VOLUME 77

SEPARATE No. 91

PROCEEDINGS

AMERICAN SOCIETY
OF
CIVIL ENGINEERS

OCTOBER, 1951



CONSUMPTIVE USE OF WATER

By Harry F. Blaney, M. ASCE

IRRIGATION DIVISION

Copyright 1951 by the AMERICAN SOCIETY OF CIVIL ENGINEERS
Printed in the United States of America

Headquarters of the Society 33 W. 39th St. New York 18, N.Y.

PRICE \$0.50 PER COPY

GUIDEPOST FOR TECHNICAL READERS

"Proceedings-Separates" of value or significance to readers in various fields are here listed, for convenience, in terms of the Society's Technical Divisions. Where there seems to be an overlapping of interest between Divisions, the same Separate number may appear under more than one item.

Technical Division	Proceedings-Separate Number
Air Transport	.42, 43, 48, 52, 60, 93, 94, 95 (Discussion: D-XXVIII, D-7, D-16, D-23, D-18, D-43)
City Planning	.58, 60, 62, 64, 93, 94 (Discussion: D-16, D-23, D-43)
Construction	.43, 50, 55, 71, 92, 94 (Discussion: D-3, D-8, D-23, D-17, D-36, D-40)
Engineering Economics	.46, 47, 62, 64, 65, 68, 69, 95 (Discussion: D-2, D-19, D-27, D-30, D-36)
Engineering Mechanics	.41, 49, 51, 54, 56, 59, 61, 88, 89, 96 (Discussion: D-5, D-XXIII, D-XXV, D-18, D-24, D-33, D-34)
Highway	.43, 44, 48, 58, 70 (Discussion: D-XXVIII, D-7, D-16, D-23, D-13)
Hydraulics	.50, 55, 56, 57, 70, 71, 78, 79, 80, 83, 86, 92, 96 (Discussion: D-XXVII, D-9, D-11, D-19, D-28, D-29)
Irrigation	.46, 47, 48, 55, 56, 57, 67, 70, 71, 87, 88, 90, 91, 96 (Discussion: D-XXIII, D-3, D-7, D-11, D-19, D-25-K, D-29, D-17, D-30, D-38, D-40, D-44)
Power	.48, 55, 56, 69, 71, 88, 96 (Discussion: D-XXIII, D-2 D-3, D-7, D-11, D-19, D-25-K, D-17, D-30, D-38, D-40, D-44)
Sanitary Engineering	.55, 56, 87, 91, 96 (Discussion: D-10, D-29)
Soil Mechanics and Foundations	.43, 44, 48, 94 (Discussion: D-4, D-XXVIII, D-7, D-43, D-44)
Structural	.42, 49, 51, 53, 54, 59, 61, 66, 89 (Discussion: D-5, D-3, D-8, D-16, D-23, D-13, D-21, D-24, D-25-K, D-32, D-17, D-33, D-34, D-39, D-42)
Surveying and Mapping	.50, 52, 55, 60, 63, 65, 68
Waterways	.41, 44, 45, 50, 56, 57, 70, 71, 96 (Discussion: D-XXVII, D-9, D-8, D-19, D-27, D-28)

A constant effort is made to supply technical material to Society members, over the entire range of possible interest. Insofar as your specialty may be covered inadequately in the foregoing list, this fact is a gage of the need for your help toward improvement. Those who are planning papers for submission to "Proceedings-Separates" will expedite Division and Committee action measurably by first studying the ASCE "Guide for Development of Proceedings-Separates" as to style, content, and format. For a copy of this Manual, address the Manager, Technical Publications, ASCE, 33 W. 39th Street, New York 18, N. Y.

The Society is not responsible for any statement made or opinion expressed in its publications

Published at Prince and Lemon Streets, Lancaster, Pa., by the American Society of Civil Engineers. Editorial and General Offices at 33 West Thirty-ninth Street, New York 18, N. Y. Reprints from this publication may be made on condition that the full title of paper, name of author, page reference, and date of publication by the Society are given.

AMERICAN SOCIETY OF CIVIL ENGINEERS

Founded November 5, 1852

PAPERS

CONSUMPTIVE USE OF WATER

By Harry F. Blaney, M. ASCE

Synopsis

Consumptive use (or evapo-transpiration) includes loss of water by evaporation of moisture from the surface of the soil and loss from interception by vegetative cover and plant transpiration. This paper discusses the subject of consumptive use with special reference to definitions, methods, and results of research. The procedure for determining consumptive use of water in valley areas from climatology, irrigation, and other data is outlined. The results of experimental measurements of evapo-transpiration by agricultural crops and natural vegetation are presented briefly. The paper provides the practicing engineer the information required to estimate consumptive use in water utilization investigations.

INTRODUCTION

Knowledge concerning the various phases of the hydrologic cycle is incomplete. Consumptive use of water (or evapo-transpiration) is one of the important elements in this cycle of water movement from the time water falls on the land as rain or snow until it reaches the ocean. The subject of consumptive use includes evaporation of moisture from land and transpiration by vegetation and is increasingly significant, particularly in the irrigated areas of the western part of the United States. It involves problems of water supply both surface and underground, as well as those of the management of, and general economics of, irrigation and multiple-purpose projects. Data on the use of water are essential in planning federal, state, and private irrigation projects. The consumptive-use requirement for water has become an important factor in the arbitration of controversies regarding major stream systems, such as the Rio Grande and the Colorado River, in which the welfare

Note.—Written comments are invited for publication; the last discussion should be submitted by April 1, 1952.

¹ Senior Irrigation Engr., Div. of Irrigation and Water Conservation, Soil Conservation Service, Research, U. S. Dept. of Agriculture, Los Angeles, Calif.

of the people of valleys, cities, states, and even nations is involved.^{2,3} Before the available water resources and an equitable division of the use of the waters of a drainage basin in arid and semiarid regions can be satisfactorily ascertained, careful consideration must be given to the consumptive-use requirements for water in various subbasins.^{3,4}

In water-supply and irrigation investigations, engineers are called upon to make, within a limited time, estimates of probable past, present, and future evaporation and evapo-transpiration losses from areas in river valleys. Sometimes few long-period hydrologic records, except climatological data, are available.³ The results of research on evapo-transpiration provide the practicing engineer with the basic data and methods required to estimate consumptive use in utilization of water investigations.⁴ A progressive improvement of methods and rapid increase in available literature on the subject sometimes make it difficult and tedious for the engineer who does not specialize in this field to evaluate the data properly and select the methods best suited for solving his particular problem.

DEFINITIONS AND SOME INTERPRETATIONS

Definition of Consumptive Use.—In some instances there has been confusion as to the meaning of terms employed in reports on the use of water. Such terms as consumptive use, evapo-transpiration, transpiration, total evaporation, stream-flow depletion, water requirements, irrigation requirements, and duty of water have been interchanged. In this paper the terms "consumptive use" and "evapo-transpiration" are considered synonymous. They may be defined

"The sum of the volumes of water used by the vegetative growth of a given area in transpiration and building of plant tissue and that evaporated from adjacent soil, snow, or intercepted precipitation on the area in any specified time, divided by the given area. If the unit of time is small, the consumptive use is expressed in acre-inches per acre or depth in inches, whereas, if the unit of time is large, such as a crop-growing season or a 12-month period, the consumptive use is expressed as acre-feet per acre or depth in feet or inches."

This definition has been found satisfactory for use in many reports since 1935.^{2,3,4} In 1939 it was adopted with minor changes by the Committee on Irrigation of the American Society of Agricultural Engineers and is now in general use by engineers.

Other Interpretations.—Although evaporation and transpiration have been studied for more than 200 years, the term consumptive use probably was not applied to water consumption prior to 1900. Many of the writings concerning the consumptive use of water are in the form of unpublished engineering reports. The term consumptive use, as originally applied to irrigation, was defined

² "Water Utilization, Upper Rio Grande Basin," by Harry F. Blaney, Faul A. Ewing, O. W. Israelsen, Carl Rohwer, and F. C. Scobey, National Resources Committee, Washington, D. C., February, 1938, Part III.

^{3 &}quot;Consumptive Use of Water in the Irrigated Areas of Upper Colorado River Basin," by Harry F. Blaney and Wayne D. Criddle, Div. of Irrigation and Water Conservation, Soil Conservation Service, U. S. Dept. of Agriculture, Logan, Utah, April, 1949.

^{4 &}quot;Consumptive Water Use and Requirements: Report of the Participating Agencies, Pecos River," by Harry F. Blaney, Paul A. Ewing, Karl V. Morin, and Wayne D. Criddle, Joint Investigation of the National Resources Planning Board, Washington, D. C., June, 1942.

as a seasonal loss of water in acre-feet per acre irrigated. Among the first published writings dealing directly with the consumptive use, a report of an ASCE committee in 1930 is noteworthy.⁵ The committee proposed certain definitions for consumptive use of water in a basic sense, and also for the farm, the project, and the valley. It also reviewed previous estimates of consumptive use for large river systems and made reference to 24 articles, published and unpublished, dealing with net water requirements and consumptive use. One of the definitions suggested by this committee follows:

"Consumptive Use in a Basic Sense: The consumptive use, 'U', is here defined as the quantity of water, in acre-fect per cropped acre per year, absorbed by a crop and transpired or used directly in the building of plant tissue, together with that evaporated from the crop-producing land."

In a report in 1930 on rainfall penetration and consumptive use of water, consumptive use was defined as "the sum of water used by the vegetative growth of a given area in transpiration or building of plant tissue and that evaporated from the area."

In 1934, the Committee on Absorption and Transpiration, Hydrology Section, American Geophysical Union, proposed the following definition⁷:

"Consumptive Use: The quantity of water per annum used by either cropped or natural vegetation in transpiration or in the building of planttissue, together with water evaporated from the adjacent soil, snow, or from intercepted precipitation. It is sometimes termed 'evapo-transpiration'.'

In 1935 the ASCE Special Committee of the Board of Direction, on Irrigation Hydraulics, adopted the following definitions8:

"Consumptive Use: The quantity of water transpired and evaporated from a cropped area."

"Evapo-transpiration: Combined loss of water from soils by evaporation and plant transpiration."

In some reports, the term "total evaporation" is applied to the combined evaporation and transpiration loss in drainage basins.9

The handbook¹⁰ of the ASCE Committee on Hydrology of the Hydraulics Division of the American Society of Civil Engineers, published in 1949, states:

"* * * the terms 'evapo-transpiration' and 'consumptive use' have received general acceptance, these terms denote the quantity of water

^{5 &}quot;Consumptive Use of Water in Irrigation," Progress Report of the Duty of Water Committee of the Irrigation Division, Transactions, ASCE, Vol. 94, 1930, p. 1349.
6 "Rainfall Penetration and Consumptive Use of Water In Santa Ana River Valley and Coastal Plain," by Harry F. Blaney, Colin A. Taylor, and Arthur A. Young, Bulletin No. 33, Div. of Water Resources, California State Dept. of Public Works, Pomona, Calif., 1930.

^{7 &}quot;Transactions of the American Geophysical Union," National Research Council of the National Academy of Sciences, Washington, D. C., June, 1934, Part II.

^{* &}quot;Letter Symbols and Glossary for Hydraulics," by the Special Committee for Irrigation Hydraulics, ASCE, Bulletin No. 11, 1935.
* "Hydrology," Physics of the Earth Series, Vol. IX, ed. by O. E. Meinzer, McGraw-Hill Book Co., Inc., New York, N. Y., 1942, p. 314.

^{10 &}quot;Evaporation and Transpiration," in Hydrology Handbook, Manuals of Engineering Practice—No. 28, Hydrology Committee of Hydraulics Division, ASCE, 1949, Chapter 4.

transpired by plants during their growth or retained in the plant tissue, plus the moisture evaporated from the surface of the soil and the vegetation. expressed in feet or inches depth of water lost or used in a specified time."

The definitions of other terms which are sometimes confused with consumptive use are as follows:

1. Duty of water: The quantity of irrigation water applied to a given area for the purpose of maturing its crop;

2. Irrigation requirement: The quantity of water, exclusive of precipitation, that is required for crop production. It includes surface evaporation and other economically unavoidable wastes;

3. Stream-flow depletion: The amount of water that flows into a valley, or onto a particular land area, minus the amount that flows out of the valley or off from the particular land area2;

4. Transpiration: The quantity of water absorbed by the crop and transpired and used directly in the building of plant tissue, in a specified time. It does not include soil evaporation;

5. Water requirement: The quantity of water, regardless of its source, required by a crop in a given period of time, for its normal growth under field conditions. It includes surface evaporation and other economically unavoidable wastes.

RESEARCH STUDIES

Factors Affecting Consumptive Use.-Many factors operate singly or in combination to influence the amount of water consumed by plants. The effects of these factors are not necessarily constant but may fluctuate from year to year as well as from place to place. Some involve the human factor, but others are related to natural influences and the environment.

The rate of evapo-transpiration depends on climate, soil-moisture supply, vegetative cover, soils, and land management. The factors included in climate that particularly affect consumptive use are precipitation, temperature, humidity, wind movement, and length of growing season. The quantity of water transpired by plants depends upon the amount of water at their disposal, as well as on temperature and dryness of the air, wind movement, the intensity of sunlight, the stage of the development of the plant, the amount of its foliage, and the nature of its leaf.

The effect of sunshine and heat in stimulating transpiration was studied as early as 1691, according to a review of the literature by Cleveland Abbe.11 Measurements of transpiration of various kinds of plants by L. J. Briggs and H. L. Shantz indicate a close correlation between transpiration and evaporation from free water surfaces, air temperature, solar radiation, and wet-bulb depression readings.12 Scientists have made many other studies of the effects of temperature moisture and light on plant growth.13

^{11 &}quot;A First Report of the Relations Between Climates and Crops," by Cleveland Abbe, Bulletin No. 36, Weather Bureau, U. S. Dept. of Agriculture, Washington, D. C., 1905.

 ^{12 &}quot;Daily Transpiration During the Normal Growth Period and its Correlation with the Weather,"
 by L. J. Briggs and H. L. Shantz, Journal of Agricultural Research, Vol. 7, 1916.
 13 "The Water Requirements of Plants," by Lyman J. Briggs and H. L. Shantz, Bulletin No. 284,
 Bureau of Plant Industry, U. S. Dept. of Agriculture, Washington, D. C., 1913.

Studies and Results.—Research studies by federal, state, and other type agencies have been made on evaporation from soil and evapo-transpiration at various times during the past fifty years in the United States.

One of the first studies¹⁴ of evapo-transpiration losses of irrigated crops in California was made in 1903 by Irrigation Investigations, Office of Experiment Stations, United States Department of Agriculture (USDA). At various times since that date this agency, now known as the Division of Irrigation and Water Conservation of the Soil Conservation Service, has studied and measured consumptive use of water by different agricultural crops and natural vegetation in many sections of the West, in cooperation with state engineers as well as state agricultural experiment stations, and other agencies. Usually evaporation, temperature, humidity, precipitation, and wind movement were recorded at the same time. Thus, data are available in many areas for correlating consumptive use measurements with temperature and other climatological observations.

Extensive studies in the correlation of evaporation, temperature, humidity, wind movement, and evapo-transpiration were conducted by the author in 1919, at the Irrigation Field Laboratory of the USDA, located at Denver, Colo. These studies included meteorological observations as well as measurements of evaporation from various sized pans, evaporation from soil tanks, and consumptive use of water by irrigated crops grown in tanks.¹⁵

The results of these studies and other investigations in California, New Mexico, and other areas indicate the observed evaporation data may be used as a means of estimating evapo-transpiration by phreatophytes (waterloving vegetation) having access to an ample water supply when the relation of the two is known for a particular area.^{2,4} As an example, for two locations in California, for tules growing in large tanks within the confines of a swamp area, the consumptive use, with reference to evaporation from a near-by exposed

Weather Bureau pan, was 95%. The percentage varied from month to month, increasing during the summer and becoming smaller in the cooler months, but 95% was the average obtained from a two-year record.

From long-period records of evaporation, temperature, and humidity in New Mexico and Texas, together with consumptive-use measurements at Carls-

TABLE 1.—Coefficients for Consumptive Use of Water, Carlsbad, New Mexico—1940

Type of vegetation	Depth of water table	Empirical Coefficients			
	(feet)	k_w	k.		
Sacaton Sacaton Salt cedar (tamarisk)	4 2 2	0.0044 0.0063 0.0075	0.0139 0.0154 0.0216		

bad, N. Mex., empirical formulas were developed for computing evaporation and consumptive use when temperature and humidity data are available.⁴ Consideration of these results and the factors involved is shown in the ex-

14 "Evaporation Losses in Irrigation and Water Requirements of Crops," by Samuel Fortier, Bulletin No. 177, Office of Experiment Stations, U. S. Dept. of Agriculture, Washington, D. C., 1907.

^{15 &}quot;Progress Report on Evaporation and Use of Water Studies at Denver Irrigation Field Laboratory, Colorado," by Harry F. Blaney, Div. of Irrigation, U. S. Dept. of Agriculture, Los Angeles, Calif., 1920 (unpublished).

pression:

$$u = k t p (114 - h) = k c \dots (1)$$

in which u is the monthly consumptive use (or evaporation) in inches; k is the monthly empirical coefficients; t is the mean monthly temperature, ${}^{\circ}F$; p is the monthly percentage of daytime hours of the year; h is the average monthly




Fig. 1.—Observed and Calculated Curves for Use of Water, Carlsbad, New Mexico, 1940

humidity; and c = t p (114 - h) is the monthly use index (climatic factor). The formula for annual consumptive use (or evaporation) in inches is

$$U = K_a C = K_w c_w + k_s c_s \dots (2)$$

in which K_a is the empirical coefficient for the entire year; C is the use index for entire year; k_a is the empirical coefficient for winter period; k_a is the empir-

ical coefficient for growing season or frost-free period; c_w is the use index for winter season; and c_s is the use index for growing season or frost-free period.

The values of k_w and k_s may be computed from observed values of consumptive use, temperature, and humidity by the relation k = u/c.

Computed coefficients for winter and summer consumptive use of water for natural vegetation (phreatophtes) growing along the Pecos River, based on evapo-transpiration, temperature, and humidity measurements at Carlsbad for the year 1940, are shown in Table 1. The computed annual coefficient K for lake-surface evaporation is 0.0164. The summer coefficient for alfalfa, growing in an area of high water table at Albuquerque, N. Mex., in 1936, is computed to be $k_* = 0.0170$.




Fig. 1 shows the observed and calculated curves for use of water for evaporation from a Weather Bureau pan, a tank of sacaton growth, and bare soil tank at Carlsbad in 1940. An example of the results of research studies of the transpiration use of water is shown in Fig. 2 which illustrates the difference between evaporation from a lake surface and the use of water by alfalfa and citrus trees in the San Fernando Valley, Calif. 16

The results of some measurements by various investigators of consumptive use by agricultural crops and natural vegetation are shown in Tables 2 and 3, respectively.

^{16 &}quot;Progress Report on Cooperative Research Studies on Water Utilization, San Fernando Valley, California, Irrigation Season of 1950," by Harry F. Blaney and Homer J. Stockwell, Jr., Div. of Irrigation, Soil Conservation Service, U. S. Dept. of Agriculture, June, 1941 (unpublished).

TABLE 2.—Seasonal Consumptive Use of Water by Various Irrigated Crops as Determined by Field Experiments

Location	Grow	ING SEASON PERIOD	Year	Consumptive use of water	
	From	То		(inches)	
(1)	(2)	(3)	(4)	(5)	(6)
		(a) A	LFALFA		
Bonners Ferry, Idaho. Carlsbad, N. Mex. Davis, Calif. Fort Stockton, Tex. Logan, Utah. San Fernando, Calif.	May 5 April 18 April 1 April 13 May 7 April 1	September 25 November 10 September 30 November 11 October 11 October 31		24.0 38.6 30.4 40.5 25.0 37.4	Marr and Criddle ^a Blaney ^b Veihmeyer ^c Blaney and Bloodgood Pittman and Stewart ^d Blaney and Stockwell ^e
		(b)	Corn		1
Davis, Calif Logan, Utah Vernal, Utah	June 1 June 1 June 10	September 30 September 30 September 20	1902-1929 1948	12.0 25.0 19.4	Veihmeyers Pittman and Stewarts Criddle and Petersons
		(c) C	OTTON		
Bakersfield, Calif Fort Stockton, Tex Los Banos, Calif Mesa, Ariz	April 1 April 13 May 1 April 1	October 31 November 11 November 30 October 31	1927-1930 1940 1932 1935	29.2 28.9 25.5 30.9	Beckett and Dunshees Blaney and Bloodgoods Adams and Veihmeyers Harris and Hawkins
		(d) SMAL	L GRAINS		
Bonners Ferry, Idaho Logan, Utah San Luis Valley, Colo Scottsbluff, Nebr	May 5 May 10 June 1 April 20	August 5 August 10 August 31 July 25	1930-1947 1902-1929 1936 1932-1933	17.5 17.5 14.05 14.72	Marr and Criddle ^a Pittman and Stewart ^d Blaney ^b Bowen ^j
1		(e) ORCHARD C	ROPS—ORANG	ES	
Azusa, Calif	April 1 March 1 April 1	October 31 October 31 October 31	1929 1931–1934 1929	21.8 32.4 20.9	Blaney and Taylor ^k Harris and Kinnison ^l Beckett and Pillsbury ^m
	(f) ORCHARD C	ROPS-WALNU)TS	
Tustin, Calif	April 1 April 1	September 30 September 30	1928 1929	26.30 27.43	Beckett ⁿ Beckett ⁿ
*	(g) C	RCHARD CROPS	-Deciduous	TREES	
Albuquerque, N. Mex Dr.vis, Calif Wenatchee, Wash Ontario, Calif	May 1 March 1 April 15 April 1	September 31 November 30 October 22 September 30	1936 1908 1928	26.4 23.0	Blaney ^b Veihmeyer ^c Fortier ^o Blaney and Taylor ^k
	•	(h) Por	ATOES		44. 4
Ontario, Ore	May 8 May 20 April 20 June 1	September 27 September 15 August 31 September 30	1947 1902-1929 1941-1942 1936	15.0 17.9	Marr and Criddle ^a Pittman and Stewart ^a Sanford and Criddle ^p Blaney ^b

TABLE 2.—Continued

Location	Growi	NG SEASON PERIOD	Year	Consumptive use of water	Authority	
(1)	(1) From To (inches) (2) (3) (4) (5)		(6)			
		(i) Suga	R BEETS		12	
Davis, Calif	April 1 April 15 April 20	September 30 October 15 October 15	1902-1929 1932-1936	25.20 25.0 24.0	Veihmeyer ^c Pittman and Stewart ^c Bowen ^j	
10 E 41 E 21 A BERT BOX 1 A A		(j) To	MATOES			
Davis, Calif Mercedes, Tex	June 1 March 25	October 31 June 30	1933-1935 1918-1920	22.8 17.0	Veihmeyers Rockwells	
		(k) Truc	ck Crops			
Stockton, Calif Stockton, Calif	May 1 April 1	September 30 October 31	1925-1928 1925-1928	21.4 24.6	Stout ^r Stout ^r	

 a "Consumptive Use of Water Studies in Idaho," by Wayne D. Criddle and James C. Marr, Div. of Irrigation, Soil Conservation Service, U. S. Dept. of Agriculture, Boise, Idaho, 1945.
 b "Consumptive Water Use and Requirements," by Harry F. Blaney, Karl V. Morin, and Wayne D. Criddle, The Pecco River Joint Investigation Reports of the Participating Agencies, National Resources Planning Board, 1942, pp. 170-230.

"Irrigation Studies," by Frank J. Veihmeyer, Univ. of California, Berkeley, Calif., 1939 (unpublished).

d "Twenty-Eight Years of Irrigation Experiments near Logan, Utah," by D. W. Pittman and George Stewart, Bulletin 219, Utah Agricultural Experiment Station, 1930, pp. 1-15.

"Progress Reports on Cooperative Research Studies of Water Utilization, San Fernando Valley, Calif.," by Harry F. Blaney and Homer J. Stockwell, 1940-41 (unpublished).

f "Consumptive Water Use and Requirements: A Progress Report on Colorado River Area Investiga-tions in Utah," by Wayne D. Criddle and Dean F. Peterson, Jr., Div. of Irrigation and Utah Agricultural Experiment Station, Soil Conservation Service, U. S. Dept. of Agriculture, 1949.

9 "Water Requirements of Cotton on Sandy Loam Soils in Southern San Joaquin Valley," by S. H. Beckett and Carroll F. Dunshee, Bulletin 537, California Agricultural Experiment Station, 1932, pp. 1-48.

h "Cotton Irrigation Investigations in San Joaquin Valley, California, 1926-1935," by Frank Adams, F. J. Veihmeyer, and Lloyd N. Brown, Bulletin 668, California Agricultural Experiment Station, 1942.

"Irrigation Requirements of Cotton on Clay Loam Soils in the Salt River Valley," by Karl Harris and R. S. Hawkins, *Bulletin 181*, Arizona Agricultural Experiment Station, 1942, pp. 421-459.

j "Uses and Efficiencies of Water by Some Farm Crops under Irrigation in Western Nebraska," by Leslie Bowen, Div. of Irrigation, Soil Conservation Service, U. S. Dept. of Agriculture, Scottsbluff, Nebr.,

k "Rainfall Penetration and Consumptive Use of Water in Santa Ana River Valley and Coastal Plain," by Harry F. Blaney, C. A. Taylor, and A. A. Young, Bulletin 33, Div. of Water Resources, California State Dept. of Public Works, 1930, pp. 1-162.

¹ "Use of Water by Washington Navel Oranges and Marsh Grapefruit Trees in Salt River Valley, ona," by Karl Harris, A. F. Kinnison, and O. W. Albertt, Bulletin 153, Arizona Agricultural Experiment Station, 1936, pp. 441-496.

m "Irrigation Requirements of California Crops," by Arthur A. Young, Bulletin 51, Div. of Water Resources, California State Dept. of Public Works, 1945, pp. 1-132.

"Irrigation Requirements in Southern California," by S. H. Beckett, Bulletin 32, Div. of Water Resources, California State Dept. of Public Works, 1930, Chapter IX, pp. 57-60.

"Orchard Irrigation," by Samuel Fortier, Farmers' Bulletin 1518, U. S. Dept. of Agriculture, 1927, pp. 1-27.

P Studies (unpublished) of Hollis Sanford, and Wayne D. Criddle, 1941-1943.

^q "Duty of Water in Irrigation," by William L. Rockwell, Div. of Irrigation and Texas Board of Water Engineers, Bureau of Public Roads, U. S. Dept. of Agriculture, 1914–1920.

"Consumptive Use of Water in the Delta of Sacramento and San Joaquin Rivers," by O. V. P. Stout, Bulletin 27, Div. of Water Resources, Calif. State Dept. of Public Works, 1931, pp. 68-75.

METHODS OF DETERMINING WATER USE

Scientists have tried various means to determine the amount of water transpired by plants.⁹ One of the earliest methods consisted of weighing freshly cut leaves, twigs, or stalks of growing plants immediately after cutting and at short intervals thereafter, during a period of hours or days, or until wilting commenced. Experimenters have also grown thousands of individual plants in small pots, weighing them periodically to determine the transpiration

TABLE 3.—Annual or Seasonal Consumptive Use of Water by Native Vegetation as Determined by Tanks and Soil Moisture Studies in Valley Areas

		Depth		PERIOD		Con- sump-	
Location	Vegetation	water table (inches)	From	То	Months	tive use (inches)	Authority
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
		(a)	NATIVE G	RASS			
San Bernardino, Calif		a	Oct. 1928	Sept. 1929	12	10.0	Blaney and Taylorb
Ontario, Calif		a	Oct. 1927	Sept. 1928	12	13.4	Blaney and Taylorb
Anaheim, Calif		a	Oct. 1927	Sept. 1928	12	12.6	Blaney and Taylorb
Cucamonga, Calif	****	a	Oct. 1929	Sept. 1930	12	15.0	Blaney and Taylorb
		(b)	NATIVE BI	LUSH			
San Bernardino, Calif		a	Oct. 1927	Sept. 1930	12	21.6	Blaney and Taylor
Muscoy, Calif			Oct. 1927	Sept. 1928	12	17.7	Blaney and Taylorb
Palmer Canyon, Calif		a	Oct. 1929	Sept. 1930	12	19.6	Blaney and Taylorb
		(c)	SALT GRA	188			
Owens Valley, Calif		18	Jan. 1911	Dec. 1911	12	48.8	Leec
Owens Valley, Calif		35	Jan. 1911	Dec. 1911	12	40.2	Leec
wens Valley, Calif		59	Jan. 1911	Dec. 1911	12	13.4	Leec
anta Ana, Calif		12 24	May 1931 May 1931	Apr. 1932 Apr. 1932	12 12	42.8 35.3	Blaney and Taylor ^d Blaney and Taylor ^d
Santa Ana, Calif		36	June 1931	Apr. 1932 Apr. 1932	11	23.8	Blaney and Taylor
Santa Ana, Calif		48	May 1931	Apr. 1932	12	13.4	Blaney and Taylor
an Luis Valley, Colo		14	Apr. 1928	Oct. 1928	7	17.8	Tipton and Harte
an Luis Valley, Colo		24	Apr. 1928	Oct. 1928	7 7	15.6	Tipton and Harte
os Griegos, N. Mex		14	Oct. 1926	Sept. 1927	12	33.2	Elder
os Griegos, N. Mex		26	Oct. 1927	Sept. 1928	12	22.7	Elder
sleta, N. Mex		8 14	June 1936	May 1937 June 1937	12 12	31.5 39.8	Blaney and Morine Blaneye
Mesilla Valley, N. Mex. Escalante Valley, Utah.		23	July 1936 May 1927	Oct. 1927	6	22.6	White/
Escalante Valley, Utah.		26	May 1927	Oct. 1927	6	17.9	White/
			(d) TREES	1			
an Luis Rey, Calif	Cottonwood, willows,	48	Apr. 1941	Mar. 1943	12	62.5	Muckel and Blaneye
afford Valley, Ariz	grass Cottonwood, willows,	84	Oct. 1943	Sept. 1944	12	91.7	Gatewood and Robinson
afford Valley, Ariz	grass Tamarisk	83.6	Oct. 1943	Sept. 1944	12	87.8	Gatewood and Robinson
afford Valley, Ariz	Tamarisk	- 24	May 1940	Dec. 1940	8 .	61.1	Turner and Halpennyi
	Tamarisk	48	May 1940	Dec. 1940	8	47.9	Turner and Halpenny
arlsbad, N. Mex	Tamarisk	36	Jan. 1940	Dec. 1940	12	57.3	Blaney and Morin
	Mesquite	120	Oct. 1943	Sept. 1944	12	32.5	Gatewood and Robinson

		Depth		PERIOD		Con- sump-	
Location	Vegetation	water table (inches)	From	tive		Authority	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)

(e) MISCELLANEOUS VEGETATION

Safford Valley, Ariz Baccharis	24	May 1940	Dec. 1940	8	52.0	Turner and Halpenny
Safford Valley, Ariz Baccharis	48	May 1940	Dec. 1940	8	39.7	Turner and Halpennyi
Safford Valley, Ariz Baccharis	72	Oct. 1943	Sept. 1944	12	54.2	Gatewood and Robinson
Escalante Valley, Utah. Grease wood	26	May 1927	Oct. 1927	6	25.2	White ^f
San Luis Valley, Colo Meadow grass	0	June 1936	Nov. 1936	6	36.3	Blanev and Morini
Carlsbad, N. Mex Sacaton	36	Jan. 1940	Dec. 1940	12	44.8	Blaney and Morini
San Bernardino, Calif Bermuda grass	24	May 1929	Apr. 1931	12	34.4	Blaney and Youngd
Fort Collins, Colo Sedge grass	6	May 1930	Oct. 1930	6	60.2	Parshall*
Fort Collins, Colo Sedge grass	18	May 1930	Oct. 1930	6	53.6	Parshall k
Isleta, N. Mex Sedge grass	3	June 1936	May 1937	12	76.9	Blaney and Morin!
Santa Ana. Calif Wire rush	24	Aug. 1930	July 1931	12	78.9	Blaney and Youngd
Fort Collins, Colo Rushes		July 1931	Oct. 1931	4	52.6	Parshall k
Victorville, Calif Tules	0		Dec. 1932	12	78.4	Blaney and Taylord
San Luis Valley, Colo Tules	0	June 1936	Nov. 1936	6	38.8	Blaneys

*Determined by Soil Sampling. No high water table. b"Rainfall Penetration and Consumptive Use of Water in Santa Ana River Valley and Coastal Plain," by H. F. Blaney, Colin A. Taylor, and Arthur A. Young, Bulletin No. 33, Div. of Water Resources, California State Dept. of Public Works, Pomona, Calif., 1930. "Hydrology," Physics of the Earth Series, ed. by O. E. Meinzer, McGraw-Hill Book Co., Inc., New York, N. Y., 1942, p. 295. "Hydrology," Physics of the Earth Series, ed. by O. E. Meinzer, McGraw-Hill Book Co., Inc., New York, N. Y., 1942, p. 295. "Water Losses Under Natural Conditions from Wet Areas in Southern California." by Harry F. Blaney, C. A. Taylor, M. G. Niekle, and A. A. Young, Bulletin No. 44, Div. of Water Resources, California State Dept. of Public Works, Pomona, Calif., 1933. "Water Utilization, Upper Rio Grande Basin," by Harry F. Blaney, Paul A. Ewing, O. W. Israelsen, Carl Rohrer, and F. C. Scobey, National Resources Committee, Washington, D. C., February, 1938, Part III. "A Method of Estimating Ground-Water Supplies Based on Discharge by Plants and Evaporation from Soil," Results of Investigations in Escalante Valley, Utah, by W. N. White, Water-Supply Paper 659-A, Geological Survey, U. S. Dept. of the Interior, Washington, D. C., 1932. "Utilization of the Waters of Lower San Luis Rey Valley, San Diego County, California," by Dean C. Muckel and Harry F. Blaney, Div. of Irrigation and Water Conservation, Soil Conservation Service, U. S. Dept. of Agriculture, Los Angeles, Calif., April, 1945. h"Use of Water by Bottom-Land Vegetation in Lower Safford Valley, Arizona," by J. S. Gatewood, T. W. Robinson, B. R. Colby, J. D. Hem, and L. C. Halpenny, Water-Supply Paper No. 1103, Geological Survey, U. S. Dept. of the Interior, Washington, D. C., 1950. "Ground-Water Inventory in the Upper Gila River Valley, New Mexico and Arizonas: Scope of Investigations and Methods Used," by S. F. Turner and L. C. Halpenny, Transactions, Am. Geophysical Union, 1941, Part III, Washington, D. C., August, 1941, p. 738.

or evapo-transpiration losses. Most of the methods of this type are artificial, and it is questionable whether the results can be extended to field conditions.

Various methods have been used by engineers to determine consumptive use of water by agricultural crops and natural vegetation under field conditions.¹⁷ Regardless of the method, the problems encountered are numerous. The source of water used by plant life, whether from precipitation alone or irrigation or ground water plus precipitation, is a factor in selecting a method. The methods most widely used in engineering investigations are:

- (a) soil-moisture studies on plots;
- (b) tank or lysimeter experiments;
- (c) ground-water fluctuations;
- (d) inflow-outflow measurements;
- (e) integration;

¹⁷ "Field Methods of Determining Consumptive Use of Water," by Harry F. Blaney, Div. of Irrigation, U. S. Dept. of Agriculture, Los Angeles, Calif., 1938.

(f) effective heat;

(g) correlation of water use, climatological data, and other data; and

(h) analysis of irrigation and precipitation records.

Soil-Moisture Studies on Plots.—This method is usually employed to determine the consumptive use of irrigated field plots in which the soil is fairly uniform and the depth to ground water is such that it will not influence the soil-moisture fluctuations within the root zone. Soil samples are taken by means of a standard soil tube before and after each irrigation, with some samples between irrigations. The samples are in 6-in. sections for the first foot and thereafter in 1-ft sections in the major root zone. Usually a great number of soil samples must be taken. 6,18

Standard laboratory practices are used in determining the moisture content of the soil samples. The samples are weighed and dried in an electric oven at 110° C and the dry weights determined. The water content of a sample is expressed as a percentage of the oven-dry weight of the soil. From the moisture percentage thus obtained, the quantity of water in acre-inches per acre (inches) removed by evaporation and transpiration from each foot of soil is computed by using the formula

$$D = \frac{P \ V \ d}{100}.$$
 (3)

in which P represents the moisture percentage by weight; V, the apparent specific gravity (or volume weight) of the soil; d, the depth of soil in inches; and D, the equivalent depth of water in inches lost by the soil (acre-inches per acre).

The total number of acre-inches of water extracted from the soil by evapotranspiration is computed for each period and later reduced to equivalent losses in acre-inches or inches for a thirty-day period. The thirty-day period losses may be plotted and a use-of-water curve for the season obtained. The average use of water for each month is taken directly from the curve (see Fig. 1). Examples of monthly consumptive-use-of-water determination made by this method are shown in Table 4 for several crops.

Tank and Lysimeter Experiments.—One of the more common methods of determining use of water by individual agricultural crops and natural vegetation is to grow the plants in tanks or lysimeters and measure the quantity of water necessary to maintain the growth satisfactorily. Tanks as large as 10 ft in diameter and 10 ft deep have been used. However, in most consumptive-use studies, the tanks are about 2 to 3 ft in diameter and 6 ft deep.6,19

The practicability of determining consumptive use by means of tanks or lysimeters is dependent on the accuracy of reproduction of natural conditions. In addition to environment, artificial conditions are the result of the limitations of soil, size of tank, or regulation of water supply. Weighing is the precise means of determining the consumptive use from tanks and this method was

¹⁸ "Irrigation Water Requirement Studies of Citrus and Avocado Trees in San Diego County, California, 1926 and 1927," by S. H. Beckett, Harry F. Blaney, and C. A. Taylor, Bulletin No. 489, College of Agriculture, Agricultural Experiment Station, Univ. of California, Berkeley, Calif., 1930.
¹⁹ "Use of Water by Native Vegetation," by Arthur A. Young and Harry F. Blaney, Bulletin No. 50, California State Div. of Water Resources, Sacramento, Calif., 1942.

used as early as 1907. However, conditions and facilities will not always permit the weighing of tanks. It has been found that all tank vegetation must be protected from the elements by a surrounding growth of the same species. Soil tanks equipped with suitable supply tanks have been used successfully in evapo-transpiration measurements from water tables at various depths. This equipment was recommended by the National Resources Committee. The tanks best adapted for this use are the double-type tank with an annular space between the inner and outer shells.^{6,19}

Charles Hamilton Lee, M. ASCE, in tests conducted in Owens Valley, Calif., 1911, employed tanks to measure the use of water by natural vegetation. Since 1919 this method has been employed in California, Colorado, and New Mexico. 2.4.6

TABLE 4.—Monthly Consumptive Use of Water as Determined by Soil Moisture Studies

			Consum	PTIVE USE	OF WATER	(Inches)		
Month	Oranges ^a	Peaches	Alfalfa	Alfalfa	Cotton	Cotton	Beets	Potatoes
	(Los Angeles, Calif.)	(Ontario, Calif.)	(Los Angeles, Calif.) ^a	(Scotts- bluff, Nebr.)	(Bakers- field, Calif.)	(Mesa, Ariz.)	(Scotts- bluff, Nebr.)	(Scotts- blufl, Nebr.)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
April May	2.2 2.2 3.1	1.0 3.5 6.7	3.3 6.7 5.4	4.0 7.0	0.8 1.1 3.3	1.1 2.2 3.6	1.9	
June July	3.4 3.7	8.0 6.5	7.8 4.2	7.1 6.4	5.6 7.7	6.0 8.8	5.2	3.4 5.8
August September. October	3.1 2.9	2.7 1.4	5.6 4.4	0.4	6.4 3.6	7.2 2.9	6.9 5.8 1.1	4.4
Total	20.6	29.8	37.4		28.5	31.8	24.2	

^a "Progress Report on Cooperative Research Studies on Water Utilization, San Fernando Valley, California, Irrigation Season of 1940." by Harry F. Blaney and Homer J. Stockwell, Jr., Div. of Irrigation, Soil Conservation Service, U. S. Dept. of Agriculture, June, 1941 (unpublished). ^b "Rainfall Penetration and Consumptive Use of Water in Santa Ana River Valley and Coastal Plain," by Harry F. Blaney, Colin A. Taylor, and Arthur A. Young, Bulletin No. 33, Div. of Water Resources, California State Dept. of Public Works, Pomona, Calif., 1930. ^c "Irrigation Studies in Arizona," by Karl Harris, Div. of Irrigation and Water Conservation, Soil Conservation Service, U. S. Dept. of Agriculture (unpublished).

Ground-Water Fluctuations.—Evapo-transpiration losses are indicated by the daily rise and fall of the water table. Diurnal measurements of the ground-water table fluctuations provide a basis for computing consumptive use of water by overlying vegetation. This method has been used successfully by the Geological Survey (USGS), U. S. Dept. of the Interior, in Escalante Valley, Utah; Santa Ana Valley, Calif.; Gila Valley, Ariz.; and other areas.^{20,21,22} The procedure used is to install observation wells, equipped with water-stage

²⁰ "A Method of Estimating Ground-Water Supplies Based on Discharge by Plants and Evaporation from Soil," Results of Investigations in Escalante Valley, Utah, by W. N. White, Water-Supply Paper 659-A, Geological Survey, U. S. Dept. of Interior, Washington, D. C., 1932.

n "Water Losses under Natural Conditions from Wet Areas in Southern California," Part I, by Harry F. Blaney, C. A. Taylor, M. G. Nickle, and A. A. Young; Part II, by Harold Troxell, Bulletin No. 44, Div. of Water Resources, California State Dept. of Public Works, Pomona, Calif., 1933.

²² "Use of Water by Bottom-Land Vegetation in Lower Safford Valley, Arizona," by J. S. Gatewood, T. W. Robinson, B. R. Colby, J. D. Hem, and L. C. Halpenny, Water-Supply Paper 1103, Geological Survey, U. S. Dept. of Interior, Washington, D. C., 1950.

registers, at representative locations in the area under consideration and to obtain graphic records of the ground-water fluctuations. The specific yield of the soil is determined by standard methods. The diurnal cycle, as indicated by the ground-water table, represents a curve showing the relation of consumptive use to ground-water discharge or recharge.

Inflow-Outflow Measurements.—Consumptive use of water by cottonwoods, willows, tules, and other riparian vegetation common to small streams may be determined by measuring the inflow to and the outflow from a selected area by means of flow recorders located on bed-rock controls. 6.21 In investigations conducted in the Upper Rio Grande, 2 the Pecos River and other river valleys, 23 engineers have used the inflow-outflow method to determine the consumptive use in irrigated valleys containing land areas up to 400,000 acres.

In the Upper Rio Grande Valley investigation the inflow-outflow method was considered to include the measurement of the amount of water flowing into and out of a given area, including precipitation and change in ground-water storage. Thus, valley consumptive use (U) is equal to the amount of water that flows into the valley during a 12-month year (I), plus the yearly precipitation on the valley floor (R), plus the water in ground storage at the beginning of the year (G_s) , minus the amount of water in ground storage at the end of the year (G_s) , minus the yearly outflow (O). All amounts are measured in acre-feet. This relation may be expressed by the equation

$$U = (I + R) + (G_s - G_s) - 0 \dots (4)$$

The difference between the storage of capillary water at the beginning and end of the year is considered negligible. It is also assumed that stream measurements are made on bed-rock controls and there is little or no subsurface flow. Example of inflow-outflow measurement in several typical areas are given in Table 5.

Integration Method.—Briefly stated, the integration method determines consumptive use by the summation of the products of consumptive use for each crop, times its area, plus the consumptive use of natural vegetation times its area, plus water surface evaporation times water surface area, plus evaporation from bare land times its area. This method was used in the Upper Rio Grande Basin² and the Pecos River⁴ investigation to determine valley consumptive use.

Before the integration method can be used successfully, it is necessary to know unit evapo-transpiration, or consumptive use of water, and the areas of various classes of agricultural crops, natural vegetation, bare land, and water surfaces. Unit values of the evapo-transpiration can be obtained by soilmoisture, tank, or some of the other methods previously described. By using aerial maps and field surveys, the areas of the various types or vegetative cover and the land and water surface areas can be determined, and the valley consumptive use computed.^{2,4}

Effective Heat.—Many formulas have been developed for determining evaporation from meteorological observations, but formulas for estimating

²³ "Consumptive Use of Water for Agriculture," by Robert L. Lowry, Jr., and Arthur F. Johnson, Transactions, ASCE, Vol. 107, 1942, p. 1243.

consumptive use are not so numerous. A few methods for determining consumptive use, based on climatic factors, have been suggested and have been found to give reasonable results. For many years irrigation engineers have used temperature data in estimating valley consumptive use of water in areas of the West.⁵ C. R. Hedke developed the effective heat method on the basis of studies of the Rio Grande. By this method consumptive use is estimated from a study of the heat units available to the crops of a particular valley. It assumes that there is a linear relation between the amount of water consumed and the quantity of available heat. Studies of the Bureau of Reclamation (USBR), United States Department of the Interior, ²³ in 1941 developed a somewhat similar method that has been employed by the USBR in making its estimates of consumptive use. This method assumes a linear relation between consumptive use and accumulated daily maximum temperatures above 32° F during the growing season.

TABLE 5.—Examples of Valley Consumptive Use of Water Determinations by Inflow-Outflow Method

. 1	Location	Year	Area	Annual Co Us		Authority	
		1001	(acres)	Total (acre-feet)	Average (feet)		
San Lu	is Valley, Colo	1925-1935	400,000	664,900	1.66	Blaney-Rohwera	
	uis Valley, Colo	1936	400,000	685,423	1.71	Blaney-Rohwera	
	uis Valley, Colo		17,300	26,215	1.52	Tipton-Harta	
	Belen, N. Mex		17,500	38,700	2.28	Blaney-Morina	
	a Valley, N. Mex		109,000	297,756	2.73	Blaney-Israelsena	
	a Valley, N. Mex		110,418	303,683	2.75	Blaney-Israelsena	
	ad, N. Mex		51,700	129,752	2.51	Blaney-Morin-Criddle	
	ad, N. Mex		51,700	119,898	2.33	Blaney-Morin-Criddleb	
New F	ork, Wyo	1939-1940	25,000		1.59	Lowry-Johnson ^c	
	gan-Illinois-Colo		43,000		1.50	Lowry-Johnson ^c	
Uncom	pahgre, Colo	1938-1940	137,700	1	2.28	Lowry-Johnson ^e	

a "Water Utilization, Upper Rio Grande Basin," by Harry F, Blaney, Paul A. Ewing, O. W. Israelsen, Carl Rohwer, and F. C. Scobey, National Resources Committee, Washington, D. C., February, 1938, Part III. b "Consumptive Water Use and Requirements: Report of the Participating Agencies, Pecos River," by Harry F. Blaney, Paul A. Ewing, Karl V. Morin, and Wayne D. Criddle, Joint Investigation of the National Resources Planning Board, Washington, D. C., June, 1942. a "Consumptive Use of Water for Agriculture," by Robert L. Lowry, Jr., and Arthur F. Johnson, Transactions, ASCE, Vol. 107, 1942, p. 1243.

In 1940, in connection with the Pecos River Joint Investigation,⁴ a method of estimating consumptive use by natural ground-water vegetation and by irrigated crops having access to an ample water supply was developed. Correlation of the relationship between evapo-transpiration, evaporation, temperature, percent of daytime hours, length of growing season, and humidity was used in the estimate.²⁴ In 1945 this method was modified to make possible estimates of consumptive use for areas in which humidity records are not available.²⁵ This method has been used by the Soil Conservation Service and other agencies in estimating rates of water consumption in many areas.

²⁴ "Evaporation and Consumptive Use of Water Empirical Formulas," by Harry F. Blaney and Karl V. Morin, *Transactions*, Part I, Am. Geophysical Union, August, 1942, p. 76.

^{25 &}quot;A Method of Estimating Water Requirements in Irrigated Areas from Climatological Data," by Harry F. Blaney and Wayne D. Criddle, Div. of Irrigation and Water Conservation, Soil Conservation Service, U. S. Dept. of Agriculture, Logan, Utah, 1945.

CORRELATION OF WATER-USE, CLIMATOLOGICAL DATA AND OTHER DATA

Actual measurements of consumptive use under each of the physical and climatical conditions of any large area are expensive and time consuming. Therefore, some rapid method of transferring the results of careful measurements, made in several areas, to other areas of similar conditions is needed. The Division of Irrigation and Water Conservation, Soil Conservation Service, has developed such a method to estimate rates of water consumption. ^{26,27} Briefly, the procedure is to correlate existing consumptive-use data with monthly temperature, percentage of daytime hours, precipitation, frost-free (growing) period, or irrigation season. The coefficients so developed for different crops are used to translocate or transpose consumptive-use data from one section to other areas in which climatological data alone are available.

TABLE 6.—Tentative Normal Consumptive-Use Coefficients for the More Important Irrigated Crops and Natural Vegetation of the West

Item	Length of growing season or period	Consumptive-use coefficient (K)
(a) IRRIGATED LAND		
Alfalfa Beans Corn. Cotron. Citrus orchard Deciduous orchard Pasture, grass, hay, annuals. Potatoes Rice Small grains Sorghum Sugar beets	frost-free 3 months 4 months 7 months 7 months frost-free frost-free 3 months 3 to 4 months 5 months 5 months	0.85 .65 .75 .62 .55 .65 .75 .70 1.00 .75
(b) NATURAL VEGETATION ^a		
Very dense (large cottonwoods, willows) Dense (tamarisk, willows) Medium (small willows, tamarisk) Light (saltgrass, sacaton)	frost-free frost-free frost-free frost-free	1.30 1.20 1.00 ^b .80

^a Ample moisture available from ground water or water table. ^b K = 1.00 for lake evaporation in arid areas and 0.90 for lake evaporation in coastal areas.

Although it is recognized that consumptive use of water is affected by numerous independent and related variables, records of temperatures and precipitation are by far the most universally available data throughout the western states. Research studies also indicate that, of all the climatic factors, these elements, together with daylight, undoubtedly have the greatest ininfluence on plant growth. Disregarding the unmeasured factors, consumptive use varies with the temperature and the extent of daytime hours, and with the available moisture (precipitation, irrigation, and ground water). By multiply-

^{28 &}quot;Irrigation Practice and Consumptive Use of Water in Salinas Valley, California," by Harry F. Blaney and Paul A. Ewing, Div. of Irrigation and Water Conservation, Soil Conservation Service, U. S. Dept. of Agriculture, Los Angeles, Calif., June, 1946.

²⁷ "Determining Water Requirements in Ir. igated Areas from Climatological and Irrigation Data," by Harry F. Blaney and Wayne D. Criddle, Div. of Irrigation and Water Conservation, Soil Conservation Service, U. S. Dept. of Agriculture, SCS-TP-96, Washington, D. C., 1950.

ing the mean monthly temperature (t) by the monthly percentage of daytime hours of the year (p), a monthly consumptive-use factor (f) is obtained. Then it is assumed that the consumptive use varies directly as this factor when an ample water supply is available. Expressed mathematically

$$U = K F = \sum k f \dots (5)$$

in which U is the consumptive use of crop (or evaporation) in inches for any period; F is the sum of the monthly consumptive-use factors for the period (sum of the products of mean monthly temperature and monthly percentage of daytime hours of the year); K is the empirical consumptive-use coefficient (irrigation season or growing period); and the other terms are as previously defined.




Fig. 3.—Relation of Consumptive Use to Temperature and Percentage of Daylight Hours

The consumptive-use factor (F) for any period may be computed for areas in which monthly temperature records are available. Then, by knowing the consumptive-use coefficient (k) for a particular crop in some locality, an estimate of the water use by the same crop in some other area may be made by application of Eq. 5.

The consumptive-use coefficients (K) for the more important irrigated crops (grown under normal conditions in the West) and natural vegetation are shown in Table 6. These coefficients were developed from actual measurements of consumptive use in tank and soil-moisture field studies; and inflow-

outflow measurements made throughout the West over a period of years by the Division of Irrigation and Water Conservation and other agencies. These coefficients are based on the assumption that the plants or trees receive a full water supply throughout the growing season or frost-free period. Fig. 3

TABLE 7.—Monthly Percentage of Daytime Hours of the Year for Latitudes 24° to 50° North of the Equator

Month				1	Latitui	DES IN	Degree	s Nort	TH OF E	QUATOR				
Month	24	26	28	30	32	34	36	38	40	42	44	46	48	50
January	7.58	7.49	7.40	7.30	7.20	7.10	6.99	6.87	6.76	6.62	6.49	6.33	6.17	5.98
February	7.17	7.12	7.07	7.03	6.97	6.91	6.86	6.79	6.73	6,65	6.58	6.50	6.42	6.32
March	8.40	8.40	8.39	8.38	8.37	8.36	8.35	8.34	8.33	8.31	8.30	8.29	8.27	8.25
April	8.60	8.64	8.68	8.72	8.75	8.80	8.85	8.90	8.95	9.00	9.05	9.12	9.18	9.25
May		9.38	9.46	9.53	9.63	9.72	9.81	9.92	10.02	10.14	10.26	10.39	10.53	10.69
June	9.20	9.30	9.38	9.49	9.60	9.70	9.83	9.95	10.08	10.21	10.38	10.54	10.71	10.93
July	9.41	9.49	9.58	9.67	9.77	9.88	9.99	10.10	10.22	10.35	10.49	10.64	10.80	10.99
August	9.05	9.10	9.16	9.22	9.28	9.33	9.40	9.47	9.54	9.62	9.70	9.79	9.89	10.00
September	8.31	8.31	8.32	8.34	8.34	8.36	8.36	8.38	8.38	8.40	8.41	8.42	8.44	8.44
October	8.09	8.06	8.02	7.99	7.93	7.90	7.85	7.80	7.75	7.70	7.63	7.58	7.51	7.43
November	7.43	7.36	7.27	7.19	7.11	7.02	6.92	6.82	6.72	6.62	6.49	6.36	6.22	6.07
December	7.46	7.35	7.27	7.14	7.05	6.92	6.79	6.66	6.52	6.38	6.22	6.04	5.86	5.65
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

shows the relation of consumptive use (U); consumptive-use factor (F); and coefficient (K). Monthly percentage of annual daytime hours computed from sunshine tables²⁸ are shown in Table 7.

Examples of consumptive-use rates in the Upper Colorado River Basin,³ computed by the above method, are shown in Table 8.

The consumptive-use coefficients shown in Table 6 are suitable for average conditions in the semiarid areas of the West. Studies to develop correction

TABLE 8.—Estimates of Normal Unit Consumptive-Use Rates During the Irrigation Period for Irrigated Crops for Several Localities in the Upper Colorado River Basin

Location	Normal Rate of Consumptive Use, in Inches						
	Alfalfa	Pasture	Grains	Corn			
Chinle, Arizona Eagle, Colorado Cortez, Colorado	28.3	24.2	15.4	20.1			
Eagle, Colorado	18.5	16.2	11.6	11.6			
Cortez, Colorado	24.5	21.5	14.6	18.9			
Dirango, Colorado	22.9	20.1	14.0	17.1			
	32.1	28.4	15.3	20.4			
Green River, Utah	32.5	28.6	16.2	21.7			
Price, Utah	26.7	23.5	15.4	19.9			
Eden, Wyoming	17.2	15.1	13.4				
Pinedale, Wyoming	14.8	13.0					

^a Values used in the formula U = KF are: Alfalfa, 0.85; pasture, 0.75; grain, 0.75; and corn, 0.75.

factors for humid conditions in the eastern states have been undertaken. It is anticipated that evaporation, humidity, or solar radiation records can be

²⁸ "Sunshine Tables," Bulletin No. 805, Weather Bureau, U. S. Dept. of Agriculture, Washington, D. C., 1905.

used for this purpose until measured consumptive-use data are available. Basic solar radiation data are now available at many stations throughout the United States.

ANALYSIS OF IRRIGATION AND PRECIPITATION RECORDS

In the consumptive-use studies conducted on the Pecos River in 1940, a method of determining unit consumptive use for various agricultural crops from irrigation and precipitation was developed. This method has been successfully employed in several areas in California. It consists of estimating evaporation and transpiration losses during the irrigation season and winter period based on data of the number of irrigations and depth of water applied, farm duty of water, and efficiency of irrigation water application for various crops and soils; and precipitation and evaporation from free water surface. Precipitation may be a major factor in winter consumptive use.

A field survey of conditions through the Pecos Valley area was made. Water superintendents of irrigation district and water companies, engineers, county farm advisors, and farmers were interviewed to ascertain the quantity of irrigation water supplied to various crops, both for the growing season and the winter period. Precipitation, irrigation, and other records were collected and estimates made of the efficiency of irrigation for various crops in the different areas. The latter data were supplemented by water-application efficiency studies, made on typical farms by measuring the quantity of water applied and taking soil samples before and after irrigation. By this means it was possible to determine the proportion of soil moisture available for crop transpiration use within the root zone. These data were analyzed and the unit rate of water consumption for each crop computed. The total consumptive use in acre-feet may be computed by multiplying unit values of consumptive use for each crop by its area.

ACKNOWLEDGMENT


This paper serves as an introduction to several others on this same general subject^{29,30,31,32} and should provide the background necessary for complete understanding of the specialized phases of the field.

¹⁰ "Consumptive Use of Water by Forest and Range Vegetation," by L. R. Rich, Proceedings-Separate No. 90, ASCE, October, 1951.

*2 "Consumptive Use in Municipal and Industrial Areas," by George B. Gleason, Proceedings-Separate No. 99, ASCE, November, 1951.

²⁹ "Consumptive Use in the Rio Grande Basin," by Robert L. Lowry, Proceedings-Separate No. 97, ASCE, November, 1951.

^{31 &}quot;Consumptive Use of Water on Irrigated Land," by Wayne D. Criddle, Proceedings-Separate No. 98, ASCE, November, 1951.



CURRENT PAPERS AND DISCUSSIONS

		CUI	RENT PAPERS AND DISCUSSIONS	
Proceedings- Separate Number	Da Avail		Title and Author	Discus- sion closes
57	Mar.,		"Design of Irrigation Systems," by W. H. Nalder	
58	Mar.,		"Highway Planning in Turkey," by H. E. Hilts	Aug. 1
59	Mar.,		"Limit Design of Beams and Frames," by H. J. Greenberg and W. Prager	Aug. 1
60	Mar.,	1951	"Surveying and Mapping Requirements for Modern City Plan- ning," by Charles A. Blessing "Structural Damping in Suspension Bridges," by the late Fried- rich Bleich and L. W. Teller.	.Aug. 1
61	Apr.,	1951	"Structural Damping in Suspension Bridges," by the late Friedrich Bleich and L. W. Teller	.Sept. 1
62	Apr.,	1951	"The New Towns Program in Great Britain," by T. C. Coote	Sept. 1
63	Apr.,	1951	"National Geodesy-Status and Planning," by Leo Otis Colbert	
64	May,	1951	"Planning the National Capital: Objectives and Problems of Attainment," by Ulysses S. Grant, III	.Oct. 1
65	May,	1951	"National Topographic Mapping," by W. E. Wrather	Oct. 1
66	May,	1951	"Lateral Forces of Earthquake and Wind," by a Joint Committee of the San Francisco, Calif., Section, ASCE, and the Structural Engineers Association of Northern California	.Oct. 1
67	June,	1951	"Model and Prototype Studies of Sand Traps," by Ralph L. Parshall	. Nov. 1
68	June,	1951	"International Cartography," by Robert H. Randall	. Nov. 1
69	June,	1951	"Valuation and Depreciation Related to Income Tax," by Maurice R. Scharff	
70	June,		"Regime Theory for Self-Formed Sediment-Bearing Channels," by T. Blench	. Nov. 1
71	June,		"Rectification of the Papaloapan River in Mexico," by Reynaldo Schega	
72	June,		"Design of Large Coal Bunkers," by Paul Rogers	Nov. 1
73	July,		"Health Implications of Air Pollution," by J. J. Bloomfield	. Dec. I
74	July,		"Acceptable Standards for Natural Waters Used for Bathing," by Charles R. Cox	Dec. 1
75	July,		"Base Course Drainage for Airport Pavements," by Arthur Casagrande and William L. Shannon	.Dec. 1
76	July,		"Model Tests Using Low-Velocity Air," by James W. Ball	. Dec. 1
77	Aug.,	1951	"Buckling Stresses for Flat Plates and Sections," by Elbridge Z. Stowell, George J. Heimerl, Charles Libove, and Eugene E. Lundquist.	Jan. 1
78	Aug.,	1951	"River Channel Roughness," by Hans A. Einstein and Nicholas L. Barbarossa.	Jan. 1
79	Aug.,	1951	"Stage Predictions for Flood Control Operations," by Ralph E. King.	.Jan. 1
80	Aug.,		"Mississippi River Valley Geology Relation to River Regime," by Harold N. Fisk	
81	Aug.,		"Petenwell Hydroelectric Project," by E. Montford Fucik	Jan. 1
82	Sept.,		"Pressures in a Shallow Rectangular Bin," by Raymond L. Moore and J. R. Shaw.	.Feb. 1
83	Sept.,		"Waterway Traffic on the Great Lakes," by John R. Hardin	. Feb. 1
84 85	Sept.,		"Longitudinal Mixing Measured by Radioactive Tracers," by Harold A. Thomas, Jr., and Ralph S. Archibald	.Feb. 1
86	Sept.,		Aultman. "Ground-Water Movement Controlled Through Spreading," by	.Feb. 1
87	Oct.,		Paul Baumann	.Feb. 1
88	Oct.,		"Sewage Reclamation by Spreading Basin Infiltration," by Ralph Stone and William F. Garber" "Experimental Study of Water Flow in Annular Pipes." by W. M.	.Mar. 1
89	Oct.,		"Experimental Study of Water Flow in Annular Pipes," by W. M. Owen. "Deflections in Gridworks and Slabs," by Walter W. Ewell, Shigeo Okubo, and Joel I. Abrams.	.Mar. 1
90	Nov.,	1951	"Consumptive Use of Water by Forest and Range Vegetation,"	
			by L. R. Rich	.Apr. 1
91	Nov.,		"Consumptive Use of Water," by Harry F. Blaney	.Apr. 1
92	Nov.,		"Experimental Investigation of Fire Monitors and Nozzles," by Hunter Rouse, J. W. Howe, and D. E. Metzler	.Apr. 1
93	Nov.,		"Aircraft Design as Related to Airport Standards," by Milton W. Arnold	
94	Nov.,		"Friendship International Airport," by Benjamin Everett Beavin	Apr. 1
95	Nov.,		"Directional Requirements for Airport Runways," by Ralph H. Burke and Harry Otis Wright, Jr.	.Apr. 1
96	Nov.,	1951	"Surface Curves for Steady Nonuniform Flow," by Robert B. Jansen	.Apr. 1

AMERICAN SOCIETY OF CIVIL ENGINEERS

OFFICERS FOR 1951

PRESIDENT GAIL A. HATHAWAY

VICE-PRESIDENTS

Term expires January, 1952:

FRED C. SCOBEY ALBERT HAERTLEIN

Term expires October, 1952:

WILLIAM R. GLIDDEN DANIEL V. TERRELL

DIRECTORS

WALDO G. BOWMAN HAROLD L. BLAKESLEE PAUL L. HOLLAND EDMUND FRIEDMAN S. T. HARDING

OTTO HOLDEN FRANK L. WEAVER GORDON H. BUTLER G. BROOKS EARNEST WALTER J. RYAN* GEORGE W. LAMB

Term expires January, 1952: Term expires January, 1953: Term expires October, 1952: MORRIS GOODKIND MILTON T. WILSON Term expires October, 1953:

Term expires October, 1953:
KIRBY SMITH
FRANCIS S. FRIEL
LOUIS R. HOWSON
WALLACE L. CHADWICK
NORMAN R. MOORE
BURTON G. DWYRE

PAST-PRESIDENTS

Members of the Board

FRANKLIN THOMAS

ERNEST E. HOWARD

TREASURER

EXECUTIVE SECRETARY WILLIAM N. CAREY

CHARLES E. TROUT

ASSISTANT SECRETARY

ASSISTANT TREASURER GEORGE W. BURPEE

E. L. CHANDLER

PROCEEDINGS OF THE SOCIETY

SYDNEY WILMOT Manager of Technical Publications

HAROLD T. LARSEN Editor of Technical Publications

JOHN C. REHFIELD Assoc. Editor of Technical Publications

COMMITTEE ON PUBLICATIONS

WALDO G. BOWMAN

FRANCIS S. FRIEL S. T. HARDING

OTTO HOLDEN

LOUIS R. HOWSON

NORMAN R. MOORE

^{*} Deceased.